

Suggested citation for this article: Young DS, Kramer LD, Maffei JG, Dusek RJ, Backenson PB, Mores CN, et al. Molecular epidemiology of eastern equine encephalitis virus, New York. *Emerg Infect Dis.* 2008 Mar; [Epub ahead of print]

Molecular Epidemiology of Eastern Equine Encephalitis Virus, New York

David S. Young,* Laura D. Kramer,*† Joseph G. Maffei,* Robert J. Dusek,‡
P. Bryon Backenson,*† Christopher N. Mores,§ Kristen A. Bernard,*† and Gregory D. Ebel¶

*New York State Department of Health, Albany, New York, USA; †School of Public Health, Albany, New York, USA; ‡National Wildlife Health Center, Madison, Wisconsin, USA; §University of Florida, Vero Beach, Florida, USA; and ¶University of New Mexico School of Medicine, Albuquerque, New Mexico, USA

Perpetuation, overwintering, and extinction of eastern equine encephalitis virus (EEEV) in northern foci are poorly understood. We therefore sought to describe the molecular epidemiology of EEEV in New York State during current and past epizootics. To determine whether EEEV overwinters, is periodically reintroduced, or both, we sequenced the E2 and partial NSP3 coding regions of 42 EEEV isolates from New York State and the Eastern Seaboard of the United States. Our phylogenetic analyses indicated that derived subclades tended to contain southern strains that had been isolated before genetically similar northern strains, suggesting southern to northern migration of EEEV along the Eastern Seaboard. Strong clustering among strains isolated during epizootics in New York from 2003–2005, as well as from 1974–1975, demonstrates that EEEV has overwintered in this focus. This study provides molecular evidence for the introduction of southern EEEV strains to New York, followed by local amplification, perpetuation, and overwintering.

Eastern equine encephalitis virus (EEEV; genus *Alphavirus*: family *Togaviridae*) is maintained in an enzootic cycle between ornithophilic mosquitoes and birds. The virus causes disease in some avian hosts and in incidental hosts, such as horses and humans; case-fatality rate in humans is $\approx 33\%$ (1). Virus activity has been detected in North and South America. In the United States, EEEV has been detected along the Gulf of Mexico and the Atlantic Seaboard as well as in inland foci near the Great Lakes, including upstate New York. The EEEV virion contains a single-stranded, positive-sense RNA genome of ≈ 12 kb. The 5' end of the genome

encodes 4 nonstructural proteins: NSP1, NSP2, NSP3, and NSP4. The structural proteins are encoded in the 3' third of the genome and are translated from a subgenomic RNA, 26S, resulting in 5 protein products: C, 6K, E1, E2, and E3 (2). Previous sequencing studies analyzed genetic relationships of EEEV strains in the Western Hemisphere and compared strains distributed across widespread geographic regions (3–6). EEEV has 4 distinct genetic lineages; lineage I consists of highly conserved strains from North America, and lineages II–IV encompass strains from Central and South America (3).

Outbreaks of EEEV in New York have been observed periodically since 1952, when the virus was first detected in pheasants (7). Disease in humans and/or horses has been noted on Long Island, in the lower Hudson Valley, and in central upstate New York; the last known human case in New York occurred in 1983 in Onondaga County (8). Most EEEV activity in New York has occurred in counties bordering Oneida Lake in central upstate New York (Figure 1A). Most of the activity in this region has been concentrated in the Big Bay–Toad Harbor Swamp complex in Oswego County and Cicero Swamp in Onondaga County (8). *Culiseta melanura* (Coquillett), the main enzootic vector of EEEV, breeds abundantly in these swamps (9). Localized epizootics in the counties of Oswego and Onondaga have been documented in a transmission focus during 1971–1977, 1982–1983, and 1990–1991 (8,10–13) and from 2003 to the present (2007; D.S. Young et al., unpub. data). Between these epizootic periods, EEEV was undetectable in horses and birds and only infrequently detected in mosquito pools (D.S. Young et al., unpub. data) (8). From 1992 through 1997 in upstate New York, EEEV was detected in 18 mosquito pools from Onondaga County (1994) and 3 mosquito pools from Oswego County (1996); no equine or avian cases were detected (D.S. Young et al., unpub. data). From 1998 through 2002, EEEV was not detected in mosquitoes or vertebrates in New York. However, in 2003, EEEV activity increased across New York with the emergence of the current epizootic (2003–2007) in the Onondaga and Oswego Counties region.

Patterns of localized perpetuation, overwintering, and extinction of EEEV in transmission foci are poorly understood. To determine whether EEEV overwinters locally in temperate regions such as upstate New York or whether annual reintroduction is required to reinitiate the transmission cycle, we compared nucleotide sequences comprising the entire E2 coding region and part of the NSP3 coding region. We examined 35 strains isolated in New York during 1971–1975 and 2003–2005 and 7 strains collected along the Eastern Seaboard of the United States

during 2002–2003. Using these data, we described the molecular epidemiology of EEEV strains collected during the current and past epizootics in New York.

Materials and Methods

EEEV Detection and Isolation

Isolates from New York State and the Eastern Seaboard were sequenced for this study (Appendix Table). Strains originating outside of New York were isolated from avian serum samples, which were collected during a study conducted by the US Geological Survey and stored at -80°C until inoculation onto cell culture. All EEEV strains from within New York were collected from mosquito, avian, and equine samples that were submitted to the Wadsworth Center's Arbovirus Laboratories as a part of surveillance efforts by the New York State Department of Health. EEEV strains isolated during 1971–1975 were obtained from our archives.

To obtain mosquito-derived EEEV strains, mosquitoes were collected from May through October by local county health department staff, who used standard miniature light or gravid traps. Mosquitoes were identified to the species level, and pooled samples of 10–50 mosquitoes in 2-mL microfuge tubes were submitted to the Arbovirus Laboratories for analysis. Tubes contained a steel ball-bearing (Daisy Brand, Rogers, AR, USA), and to each tube we added 1 mL of mosquito diluent (20% heat-inactivated fetal bovine serum [FBS] in Dulbecco's phosphate-buffered saline with 50 $\mu\text{g/mL}$ penicillin/streptomycin, 50 $\mu\text{g/mL}$ gentamicin, and 2.5 $\mu\text{g/mL}$ amphotericin B). Pools were homogenized by using a mixer mill (Retsch, Haan, Germany) at 24 cycles/s for 30 s and centrifuged for 4 min at 6,000 rpm at room temperature. The clarified homogenate was transferred to a new microcentrifuge tube and stored at -80°C until testing.

To obtain vertebrate-derived EEEV strains, samples of horse brains were submitted by the Wadsworth Center's Rabies Laboratory, and samples of avian kidney, heart, and brain were submitted by the New York State Department of Environmental Conservation Wildlife Pathology Unit. Each horse brain was excised into 3 separate 1–3 mm^3 sections, which were pooled for testing for each horse. Avian tissues were tested by excising the same-size portion from 1 of the tissues or by pooling sections of all 3 tissues. Excised tissues were placed in 2-mL microfuge tubes containing a ball-bearing and 1 mL of BA-1 virus diluent (M199 with Hanks'

salts and L-glutamine; [Mediatech, Herndon, VA, USA] in sterile distilled water with 0.05 M Tris(hydroxymethyl)aminomethane, 1% bovine serum albumin, 0.35 g/L sodium bicarbonate, 100 U/mL penicillin, 100 U/mL streptomycin, 1 µg/mL amphotericin B, and 20% FBS.) Hydrogen chloride was added to the diluent to bring the pH to 7.4. Samples were homogenized on the mixer mill at 24 cycles/s for 4 min and centrifuged for 5–8 min at 10,000 rpm at 4°C.

Virus was isolated by inoculating 100 mL of supernatant from mosquito pools, vertebrate tissues, or avian serum onto confluent monolayers of African green monkey kidney (Vero) cells grown in 6-well plates. Plates were incubated for 1 h at 37°C in 5% carbon dioxide, with gentle rocking every 15 min. To each well, 3 mL of maintenance medium (1× minimum essential medium with Earle's salts [Invitrogen, Carlsbad, CA, USA], 2% heat-inactivated FBS, 1% 100× L-glutamine, 0.15% sodium bicarbonate, 1% penicillin–streptomycin, 0.1% amphotericin B, and 0.1% gentamicin diluted in sterile distilled water) was added, and plates were returned to the incubator and observed daily. If cytopathic effect was observed, the infecting virus was identified by either immunofluorescence assay (14) or reverse transcription PCR (RT-PCR) by using One-Step RT-PCR kit and protocol (QIAGEN, Valencia, CA, USA) as directed by the manufacturer. Primer sequences and cycling parameters are described elsewhere (15). Target bands were examined under UV light after electrophoresis on a 1.5% agarose gel. Samples that were positive for EEEV were stored at –80°C until use.

RNA Extraction and RT-PCR

RNA was extracted from the Vero cell culture supernatant after a single passage, the original sample (no passage), or isolates with an unknown passage history by using the RNeasy kit (QIAGEN) as directed by the manufacturer. The entire E2 coding region was amplified in 2 separate reactions. To produce overlapping fragments, primers EEE8460 (5'-AGAATCCACACGAAACACTCACCA-3') and EEE9200c (5'-ATCCGTGCAGGTGGTTGTATGGTC-3') were used for the first reaction, and primers EEE9105 (5'-TCCACAGTGCCAAGGTGAAAA-3') and EEE9887c (5'-CTGCAAGTGGGATAAGCGTCTG-3') were used for the second reaction. The partial NSP3 coding region was amplified by using primers EEE4836 (5'-CAGAGCGAGTTTACAGATTACG-3') and EEE5477c (5'-AACGGCGAACGACTGAA-3'). Sample RNA (5 µL) was added to 45 µL of One-Step RT-PCR master mix (QIAGEN) prepared according to the manufacturer. Several drops of mineral oil were added on top of each reaction.

Samples were reverse transcribed for 30 min at 55°C and heat inactivated at 95°C for 5 min. To eliminate residual RNA, RNase was added to each reaction after reverse transcription. Samples were amplified by PCR according to the following thermocycler conditions: 94°C for 10 min; 39 cycles of 94°C for 30 s, 55°C for 30 s, 72°C for 45 s; and 72°C for an additional 10 min. PCR product (40 µL) was added to 4 µL of BlueJuice loading dye (Invitrogen, Carlsbad, CA, USA), and loaded onto an agarose gel containing 0.4 µg/mL of ethidium bromide. DNA underwent electrophoresis and was examined under UV light. Samples were sequenced on either an ABI 3100 or ABI 3700 automated DNA sequencer (Applied Biosystems, Foster City, CA, USA) at the Wadsworth Center Molecular Genetics Core facility. (Sequencing primers are available from the authors upon request.)

Phylogenetic Analysis

Trace files were compiled by using the SeqMan module of Lasergene (DNASTar, Madison, WI, USA), with a minimum of 2-fold base-call redundancy required for all sequences. Consensus sequences for each strain sequenced in this study and reference strains obtained from GenBank were aligned by the ClustalV method (16) in the MegAlign module of Lasergene. Two phylogenetic trees were produced for the E2 coding region analysis: the main E2 tree, which included all strains included in this study, and the E2 subset tree, which contained only strains sequenced in this study and only those sequences from GenBank for which both E2 and NSP3 sequence data were available. The main E2 tree was generated by maximum likelihood in PAUP version 4.0b10 (Sinauer Associates, Sunderland, MA, USA) by using the HKY85+G model with relevant parameters estimated from the data. The robustness of the branching pattern was estimated by performing 1,000 neighbor-joining bootstrap replicates under the maximum-likelihood substitution model, also using PAUP; these values are presented on the maximum likelihood tree. The E2 subset and NSP3 trees were generated by neighbor-joining analysis with 1,000 bootstrap replicates by using the Kimura 2-parameter model in MEGA2 (17). Nucleotide sequences of newly sequenced strains were deposited in GenBank (see online Appendix Table for accession numbers). Nucleotide diversity (π) and Tajima's D statistics were computed by using DnaSP (18).

Results

Virus Strains

We sequenced 42 EEEV strains, which represented various geographical locations, hosts, and isolation dates (Appendix Table). Of the 42 strains, 29 were from central upstate New York. Of those 29 strains, 13 were isolated during the 1970s epizootic, and the remainder were isolated during the current epizootic. The remaining strains sequenced were from various locations in the southeastern United States (Louisiana, Florida, and Virginia), New Jersey, and New York outside the central upstate focus (counties of Suffolk, Orange, Sullivan, Ulster, and Chemung) (Figure 1).

E2 Phylogenetic Analysis

Phylogenetic analysis of the E2 coding region demonstrated that all isolates sequenced in this study belonged to lineage I (Figure 2), and showed strong spatiotemporal clustering. The 3 strains isolated from Oswego County in 1971 (NY71a, NY71b, and NY71c) clustered together as a result of almost identical E2 coding regions. All 8 strains isolated from Oswego County in 1974 (NY74a–h) and the only strain isolated in 1975 (NY75) grouped together strongly and formed the Oswego74 clade (Figures 2–4). Strong clustering was also evident among strains isolated mainly from Onondaga County during 2003–2005 (the Onondaga03 clade, Figures 2–4). Of the 16 strains in this clade, 13 had identical E2 coding regions (data not shown). Both NY04g and NY04j grouped together and were isolated in close geographic proximity in Sullivan and Ulster Counties, respectively, in the lower Hudson Valley.

Sequence Diversity and Phylogenetics

Examination of π , a measure of sequence diversity, confirmed the close relationships of sequences sampled during spatiotemporally defined epizootics. The π values for the Oswego74 and Onondaga03 clades were 0.00035 and 0.00030, respectively. The π value for the entire set of US strains included in our analysis was 0.01058, $\approx 30\times$ greater than intraepizootic values. The Tajima test failed to reject neutrality in either the Oswego74 or Onondaga03 clade because of extremely low genetic diversity: only a few mutations were present in each grouping. Thus, the EEEV collected during New York epizootics is generally characterized by a high degree of sequence conservation with little genetic variation among spatially and temporally related strains.

However, spatiotemporal conservation was not absolute. NY73, isolated from a horse in Onondaga County in 1973 (NY73) was most genetically similar to NY69, isolated from a pheasant in Suffolk County on Long Island in 1969. Similarly, a strain isolated from a horse in Chemung County, New York (NY04k), fell into the Onondaga03 clade, which further demonstrated occasional relaxation of the otherwise strong time-space clustering of the strains studied. Consideration of strains from outside of New York provided additional instances in which the pattern of spatiotemporal clustering was broken. Well-supported subclades frequently contained southern progenitor strains that had been isolated years before they appeared in New York or New Jersey (Figures 2–4). Examples of this trend include the following: VA03 linked with NY03b and NY04f, GA97 linked with NJ03a and NJ03b, and FL02a linked with NY04g and NY04j. In addition, strain FL02b, isolated from an ovenbird (*Seiurus aurocapillus*) collected in Florida during 2002 was highly similar to the Onondaga03 clade.

To evaluate the possibility that analysis of different coding sequences would yield different results, we studied the NSP3 coding sequences of all strains for which sequence data were available or further sequencing was possible. GenBank sequence data for the NSP3 coding region was limited; therefore, another phylogenetic tree for the E2 coding region was produced by using a subset of lineage I strains for which the NSP3 sequences were available (Figure 3). This E2 subset tree was used for comparisons with the NSP3 tree to determine whether the trees shared similar topology. Phylogenetic analysis of the NSP3 coding region produced similar overall topologies (Figure 4); both trees recognized the major clades (Oswego74 and Onondaga03) and most of the minor subclades.

Discussion

Samples that were newly sequenced for this study were selected to include relevant arthropod and vertebrate hosts (mosquitoes, birds, and horses) and were drawn from archives and ongoing arbovirus surveillance efforts. Additional sequences used in the analyses were obtained from GenBank. Strains collected from the current (2003–2005) epizootic were sequenced from primary field-derived material when possible. This was done to minimize the likelihood that adaptation to tissue culture, which has been observed for other alphaviruses (19), could bias our results. Because our analyses did not show strong passage-history–dependent clustering, it seems

likely that the sequences from strains passed once in Vero cells are accurate representations of wild-type sequences. Further, strong sequence conservation among strains collected from taxonomically diverse hosts suggests that the source of the virus (mosquitoes or horses) was unlikely to result in sequence changes that might bias our conclusions.

Previous studies provided preliminary evidence that EEEV overwinters in upstate New York (5,20). The strongest such evidence was derived from RNA sequences and fingerprints that showed strong clustering of 11 strains collected in 1990 and 1991 (5). It has been suggested that EEEV may have been relatively isolated in upstate New York for up to several years and that the virus may have overwintered from 1990–1991 (5,20). The studies we present here document genetic conservation of strains throughout 3 transmission seasons and over 2 winters, which supports the observation that EEEV may overwinter in a relatively isolated upstate New York focus. Despite the accumulating molecular epidemiologic evidence for EEEV overwintering, the precise mechanisms are poorly understood. One potential overwintering mechanism is latent or chronic infection of wild birds. In such a scenario, springtime viral recrudescence might reinstitute the transmission cycle each new season (21,22). However, the results of a serologic survey of wild birds in upstate New York during 1986–1990 failed to support this notion, showing no consistent evidence for the current or recent infection of after-hatch year birds with EEEV early in the transmission season (23). Transovarial transmission in mosquitoes also has been hypothesized as a means for overwintering, but it has not been convincingly demonstrated naturally or experimentally (24–26). The main epizootic vector, *Cs. melanura*, overwinters in the larval stage (27), so the virus would need to perpetuate in these larvae. As an alternative, predatory birds could acquire the virus by feeding on infected prey, perhaps enabling the virus to persist through winter without mosquitoes. A previous study describing the isolation of *West Nile virus* (WNV; *Flaviviridae: Flavivirus*) from a hawk in New York in winter (28) supports this idea. However, conclusive evidence for this theory does not yet exist. Moreover, despite several ecologic studies of EEEV in upstate New York, the mode of long-term persistence in enzootic transmission cycles remains obscure. Accordingly, the molecular epidemiologic studies described here were undertaken to determine more convincingly whether EEEV overwinters locally in upstate New York.

Sequence data from the E2 and NSP3 coding regions of EEEV strains collected during 2 independent multiyear epizootics, 1 in the 1970s and 1 in the 2000s, enabled us to use a

molecular approach to examine whether EEEV overwinters in temperate regions. We observed strong spatiotemporal clustering of EEEV strains, including several strains that were identical in their E2 coding region, collected in a single focus over the course of several years. For example, 16 EEEV strains isolated during 2003–2005 form the Onondaga03 clade, which strongly suggests that the virus overwintered there. The probability that this highly conserved genotype was reintroduced in each of 3 consecutive years seems quite low. The Oswego74 clade also supports overwintering of EEEV. Collectively, these data indicate that EEEV was perpetuated locally through several winters in upstate New York during elevated epizootic activity periods, with 1 dominant genotype circulating in the focus.

The history of EEEV activity in New York suggests that transmission dynamics are not uniform and that periods of relative intensity punctuate interepizootic periods, when virus is undetectable or detectable only infrequently in mosquito pools (D.S. Young et al., unpub. data) (8,29). Phylogenetic analyses suggest that epizootics occur after reintroduction of novel EEEV genotypes from southern progenitor strains. For example, the Onondaga03 clade groups strongly with the strain FL02b, which was isolated from an ovenbird in Florida in 2002. The ovenbird resides in Florida and Central America in the winter and migrates north to Canada and the northern United States, including upstate New York, in the summer (30). The current epizootic therefore appears to be the result of introduction of a southern EEEV strain similar to FL02b in 2003. Although the results presented here cannot determine the precise mode of EEEV transport along the Eastern Seaboard (e.g., trade winds have also been suggested as a mechanism for moving infected mosquitoes [31,32]), it seems likely that migratory birds are involved in virus trafficking to at least some degree.

The E2 and NSP3 phylogenetic trees demonstrate that some subclades contain southern strains isolated years before genetically similar northern strains. Such is the case with subclades GA97, NJ03a, NJ03b and VA03, NY03b, NY04f and in the E2 trees only (FL02a, NY04g, NY04j). This pattern provides evidence for regular reintroduction of EEEV into enzootic areas of New York State and New Jersey. The subclades VA03, NY03b, NY04f suggest that northward migrating birds brought the virus from the South into New York in 2003. However, the low number of samples from the Eastern Seaboard limits definitive conclusions. For this study we included only 7 samples isolated from the Eastern Seaboard in 2002 and 2003. Sequencing

additional southern strains isolated in 2004 or later, to characterize the genetic relationships between northern and southern strains in greater depth, would be beneficial.

A defining feature of the collection of EEEV sequences analyzed here is the genetic conservation within and between epizootics. We observed values of genetic diversity in the Onondaga03 and Oswego74 clades that were surprisingly similar and very low (0.00030 and 0.00035, respectively). These values are ≈ 10 -fold lower than π observed in a sample of WNV sequences collected during 1999–2003 in Suffolk County, NY (0.00241) (33). The causes for the strikingly different patterns of genetic diversity observed in these 2 viral systems could include 1) increased movement of WNV-infected birds or mosquitoes compared with EEEV-infected birds or mosquitoes, leading to more frequent introduction of novel genotypes, 2) a higher replicase error rate, and 3) relaxed selective constraint in hosts or vectors of WNV relative to EEEV. Regardless of cause, the basic evolutionary dynamics of these RNA viruses appear to differ markedly.

Overall, our data support the previous findings of Weaver et al. (5,20) and provide new insights into the ecologic and evolutionary dynamics of an ongoing EEEV epizootic. We provide evidence that the virus is introduced from southern progenitor strains which, if they become established, overwinter in upstate New York; few new genotypes were successfully introduced into the epizootic focus. Our results also highlight the relative spatiotemporal genetic conservation of the virus. To facilitate a more detailed understanding of patterns of perpetuation and spread of this important zoonotic pathogen, future monitoring of EEEV activity should focus on sampling along the entire Eastern Seaboard.

Acknowledgments

We thank John Howard for providing strain information for 1970s EEEV isolates, Robert McLean for providing avian serum, and Ward Stone for providing avian tissues. We also thank the New York State Department of Health Rabies Laboratory for providing equine tissues and the Wadsworth Center Molecular Genetics Core for sequencing.

This study was supported in part by the National Institute of Allergy and Infectious Diseases, National Institute of Health, under contract N01-AI-25490 and CDC U50/CCU223671 (to L.D.K.).

Mr Young works at the Arbovirus Laboratories of Wadsworth Center of the New York State Department of Health. His primary research involves sequencing of mosquito-borne viruses.

References

1. Centers for Disease Control and Prevention. Eastern equine encephalitis fact sheet [cited 2006 Aug 25]. Available from <http://www.cdc.gov/ncidod/dvbid/arbor/eeefact.htm>
2. Schlesiner S, Schlesinger MJ. *Togaviridae*: the viruses and their replication. In: Fields BN, Knipe DM, Howley PM, editors. Fields virology, vol.1, 3rd ed. Philadelphia: Lippincott-Raven; 1996. p. 825–7.
3. Brault AC, Powers AM, Chavez CLV, Lopez RN, Cachón MF, Gutierrez LFL, et al. Genetic and antigenic diversity among eastern equine encephalitis viruses from North, Central, and South America. *Am J Trop Med Hyg.* 1999;61:579–86. [Medline](#)
4. Weaver SC, Hagenbaugh A, Bellew LA, Netesov SV, Volchikov VE, Chang GJ, et al. A comparison of the nucleotide sequences of eastern and western equine encephalomyelitis viruses with those of other alphaviruses and related RNA viruses. *Virology.* 1993;197:375–90. [Medline](#)
5. Weaver SC, Hagenbaugh A, Bellew LA, Gousset L, Mallampalli V, Holland JJ, et al. Evolution of alphaviruses in the eastern equine encephalomyelitis complex. *J Virol.* 1994;68:158–69. [Medline](#)
6. Weaver SC, Powers AM, Brault AC, Barrett ADT. Molecular epidemiological studies of veterinary arboviral encephalitides. *Vet J.* 1999;157:123–38. [Medline](#)
7. Beaudette FR, Holden P, Black JJ, Bivens JA, Hudson CB, Tudor DC. Equine encephalomyelitis in pheasants in 1952–53. Proceedings of the 58th Annual Meeting of the United States Livestock Sanitary Association. 1954;58:309–21.
8. Howard JJ, Morris CD, Emord DE, Grayson MA. Epizootiology of eastern equine encephalitis virus in upstate New York, USA. VII. Virus surveillance 1978–85, description of 1983 outbreak, and series conclusions. *J Med Entomol.* 1988;25:501–14. [Medline](#)
9. Morris CD, Corey ME, Emord DE, Howard JJ. Epizootiology of eastern equine encephalomyelitis virus in upstate New York, USA. I. Introduction, demography and natural environment of an endemic focus. *J Med Entomol.* 1980;17:442–52. [Medline](#)
10. Morris CD, Whitney E, Bast TF, Deibel R. An outbreak of eastern equine encephalomyelitis in upstate New York during 1971. *Am J Trop Med Hyg.* 1973;22:561–6. [Medline](#)
11. Morris CD, Caines AR, Woodall JP, Bast TF. Eastern equine encephalomyelitis in upstate New York, 1972–1974. *Am J Trop Med Hyg.* 1975;24:986–91. [Medline](#)

12. Srihongse S, Grayson MA, Morris CD, Deibel R, Duncan CS. Eastern equine encephalomyelitis in upstate New York: studies of a 1976 epizootic by a modified serologic technique, hemagglutination reduction, for rapid detection of virus infections. *Am J Trop Med Hyg.* 1978;27:1240–5. [Medline](#)
13. Howard JJ, Grayson MA, White DJ, Oliver J. Evidence for multiple foci of eastern equine encephalitis virus (*Togaviridae: Alphavirus*) in central New York State. *J Med Entomol.* 1996;33:421–32. [Medline](#)
14. Ngo KA, Maffei JG, Dupuis AP II, Kauffman EB, Backenson PB, Kramer LD. Isolation of Bunyamwera serogroup viruses (*Bunyaviridae: Orthobunyavirus*) in New York State. *J Med Entomol.* 2006;43:1004–9. [Medline](#)
15. Lambert AJ, Martin DA, Lanciotti RS. Detection of North American eastern and western equine encephalitis viruses by nucleic acid amplification assays. *J Clin Microbiol.* 2003;41:379–85. [Medline](#)
16. Higgins DG, Sharp PM. Fast and sensitive multiple sequence alignments on a microcomputer. *Comput Appl Biosci.* 1989;5:151–3. [Medline](#)
17. Kumar S, Tamura K, Jakobsen IB, Nei M. MEGA2: molecular evolutionary genetics analysis software. *Bioinformatics.* 2001;17:1244–5. [Medline](#)
18. Rozas J, Sánchez-DelBarrio JC, Messeguer X, Rozas R. DnaSP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics.* 2003;19:2496–7. [Medline](#)
19. Bernard KA, Klimstra WB, Johnston RE. Mutations in the E2 glycoprotein of Venezuelan equine encephalitis virus confer heparan sulfate interaction, low morbidity, and rapid clearance from blood of mice. *Virology.* 2000;276:93–103. [Medline](#)
20. Weaver SC, Bellew LA, Gousset L, Repik PM, Scott TW, Holland JJ. Diversity within natural populations of eastern equine encephalomyelitis virus. *Virology.* 1993;195:700–9. [Medline](#)
21. Emord DE, Morris CD. Epizootiology of eastern equine encephalomyelitis virus in upstate New York, USA. VI. Antibody prevalence in wild birds during an interepizootic period. *J Med Entomol.* 1984;21:395–404. [Medline](#)
22. Crans WJ, Caccamise DF, McNelly JR. Eastern equine encephalomyelitis virus in relation to the avian community of a coastal cedar swamp. *J Med Entomol.* 1994;31:711–28. [Medline](#)
23. Howard JJ, Oliver J, Grayson MA. Antibody response of wild birds to natural infection with alphaviruses. *J Med Entomol.* 2004;41:1090–103. [Medline](#)

24. Hayes RO, Daniels JB, Anderson KS, Parsons MA, Maxfield HK, LaMotte LC. Detection of eastern encephalitis virus and antibody in wild and domestic birds in Massachusetts. *Am J Hyg.* 1962;75:183–9. [Medline](#)
25. Morris CD, Srihongse S. An evaluation of the hypothesis of transovarial transmission of eastern equine encephalomyelitis virus by *Culiseta melanura*. *Am J Trop Med Hyg.* 1978;27:1246–50. [Medline](#)
26. Watts DM, Clark GG, Crabbs CL, Rossi CA, Olin TR, Bailey CL. Ecological evidence against vertical transmission of eastern equine encephalitis virus by mosquitoes (Diptera: Culicidae) on the Delmarva Peninsula, USA. *J Med Entomol.* 1987;24:91–8. [Medline](#)
27. Means RG. Mosquitoes of New York. Part II. Genera of Culicidae other than *Aedes* occurring in New York. Bulletin 430b. Albany (NY): New York State Museum; 1987. p. 47–8.
28. Garmendia AE, Van Kruiningen HJ, French RA, Anderson JF, Andreadis TG, Kumar A, et al. Recovery and identification of West Nile virus from a hawk in winter. *J Clin Microbiol.* 2000;38:3110–1. [Medline](#)
29. Howard JJ, Grayson MA, White DJ, Morris CD. Eastern equine encephalitis in New York State. *Journal of the Florida Mosquito Control Association.* 1994;65:1–7.
30. Alsop FJ III. Birds of North America: eastern region. 1st ed. In: Greenberg R, Hamilton J, editors. New York (NY): DK Publishing; 2001.
31. Sellers RF. Eastern equine encephalitis in Quebec and Connecticut, 1972: introduction by infected mosquitoes on the wind? *Can J Vet Res.* 1989;53:76–9. [Medline](#)
32. Sellers RF, Maarouf AR. Trajectory analysis of winds and eastern equine encephalitis in USA, 1980–5. *Epidemiol Infect.* 1990;104:329–43. [Medline](#)
33. Jerzak G, Bernard KA, Kramer LD, Ebel GD. Genetic variation in West Nile virus from naturally infected mosquitoes and birds suggests quasispecies structure and strong purifying selection. *J Gen Virol.* 2005;86:2175–83. [Medline](#)

Address for correspondence: Gregory D. Ebel, Department of Pathology, University of New Mexico School of Medicine, 1 University of New Mexico, MSC 08-4640, Albuquerque, NM 87131, USA; email: gebel@salud.unm.edu




Figure 1. A) United States map showing locations of eastern equine encephalitis virus strains sequenced in this study. New York State (NY) highlighted in blue; New Jersey (NJ), Virginia (VA), Florida (FL), Louisiana (LA) highlighted in gray. Map courtesy of www.theodora.com/maps, used with permission. B) New York counties where eastern equine encephalitis virus (EEEV) strains have been located (shaded). Dotted box indicates focus of most EEEV activity.




Figure 2. Maximum-likelihood phylogenetic tree of eastern equine encephalitis virus strains, based on the complete E2 coding sequence. Numbers at the nodes indicate bootstrap confidence estimated by 1,000 neighbor-joining replicates on the maximum-likelihood tree. The tree was rooted with lineage II (Brazil56), III (Panama86), and IV (Brazil85) strains.




Figure 3. Phylogenetic tree of subset of lineage I eastern equine encephalitis virus strains, unrooted neighbor-joining analysis of E2 coding region. Strains included are identical to those used in the NSP3 coding region analysis.




Figure 4. Phylogenetic tree of NSP3 coding region of subset of lineage I eastern equine encephalitis virus strains, unrooted neighbor-joining analysis.

Appendix Table. Eastern equine encephalitis virus strains collected during the current and past epizootics, USA

Strain Designation*	Strain	Location	Date Collected	Source	Passage History†	Accession no.	
						E2	NSP3
VA33	TenBroeck	Virginia	1933	Horse	sm12, v1	U01558	
MA38	Massachusetts	Massachusetts	1938	Human	unknown	AF159550	
LA47	Decuir	Louisiana	1947	Human	P1	U01552	
LA50	Arth-167	Louisiana	1950	Mosquito	gp2, ch2	AF159551	
Brazil56	BeAn-5122	Brazil	1956	Sentinel monkey	Sm2	AF159559	
NJ60	New Jersey 60	New Jersey	1960	Mosquito	p6, sm1	U01554	
NY69	NY69-7836	Suffolk, NY	1969	Pheasant	Unknown	EF594003	EF593961
NY71a	NY71-3465	Oswego, NY	8/23/1971	Horse	Unknown	EF594004	EF593962
NY71b	NY71-20686	Oswego, NY	8/24/1971	<i>Zonotrichia albicollis</i>	Unknown	EF594005	EF593963
NY71c	NY71-20925	Oswego, NY	8/28/1971	<i>Culex restuans</i>	Unknown	EF594006	EF593964
NY73	NY73-60104	Onondaga, NY	10/1/1973	Horse	Unknown	EF594007	EF593965
NY74a	NY74-39071	Oswego, NY	8/26/1974	Horse	Unknown	EF594008	EF593966
NY74b	NY74-39073	Oswego, NY	9/2/1974	Horse	Unknown	EF594009	EF593967
NY74c	NY74-23179	Oswego, NY	9/4/1974	<i>Melospiza melodia</i>	Unknown	EF594010	EF593968
NY74d	NY74-33620	Oswego, NY	9/5/1974	<i>Culiseta (Cs.) melanura</i>	Unknown	EF594011	EF593969
NY74e	NY74-33621	Oswego, NY	9/5/1974	<i>Aedes canadensis</i>	Unknown	EF594012	EF593970
NY74f	NY74-33630	Oswego, NY	9/12/1974	<i>Ae. canadensis</i>	Unknown	EF594013	EF593971
NY74g	NY74-33927	Oswego, NY	9/14/1974	<i>Cs. melanura</i>	Unknown	EF594014	EF593972
NY74h	NY74-39100	Oswego, NY	9/20/1974	Horse	Unknown	EF594015	EF593973
NY75	NY75-39011	Oswego, NY	7/16/1975	Horse	unknown	EF594016	EF593974
MA77	ME77132	Massachusetts	1977	Mosquito	m1, C6/36-1	U01555	
WI80	WIAn-5000	Wisconsin	1980	Horse	de2, sm1, v1	U01559	
FL82	82V-2137	Florida	1982	Mosquito	Unknown	U01034	U01034
MS83	MS-4789	Mississippi	1983	Human	rd2, sm3	AF159552	
MD85	215-85	Maryland	1985	Mosquito	bhk1	U01556	
Brazil85	436087	Brazil	1985	Mosquito	sm1, v1	AF159561	
Panama86	435731	Panama	1986	Horse	v2	AF159560	
CT90	Williams	Connecticut	1990	Horse	v1	U01557	
MD90	3067-90	Maryland	1990	Mosquito	Unpassed	U01553	
FL91	Florida91-4697	Florida	1991	Mosquito	sm1, v3, bhk2	AY705241	AY705241
GA91	PorEEE	Georgia	1991	Pig	Unknown	AF159557	
TX91	VRI-7164	Texas	1991	Horse	sm1	AF159553	
FL93	FL93-939	Florida	1993	Mosquito	v1	AF159554	
TX95	PV5-2547C	Texas	1995	Mosquito	sm1	AF159555	
FL96	FL96-14834	Florida	1996	Bird	v1	AF159556	
GA97	Georgia 97	Georgia	1997	Human	v2	AY705240	AY705240
Mexico97	97-1076	Mexico	1997	Horse	v1	AF159558	
LA02	LA02-5534	Louisiana	9/28/2002	<i>Thryothorus ludovicianus</i>	v1	EF593999	EF593957
FL02a	FL02-4341	Florida	10/7/2002	<i>Vireo griseus</i>	v1	EF593997	EF593955
FL02b	FL02-4809	Florida	10/16/2002	<i>Seiurus aurocapillus</i>	v1	EF593998	EF593956
NJ03a	NJ03-1974	New Jersey	9/24/2003	<i>Empidonax traillii/alnorum</i>	v1	EF594000	EF593958
NJ03b	NJ03-1977	New Jersey	9/24/2003	<i>Passerina cyanea</i>	v1	EF594001	EF593959
VA03	VA03-2789	Virginia	9/30/2003	<i>Vireo olivaceus</i>	v1	EF594002	EF593960
NY03a	NY03-0127	Suffolk, NY	8/26/2003	<i>Cs. melanura</i>	Unpassed	EF594017	EF593975
NY03b	NY03-2657	Orange, NY	9/2/2003	<i>Corvus brachyrhynchos</i>	Unpassed	EF594018	EF593976
NY03c	NY03-0248	Onondaga, NY	9/11/2003	<i>Cs. melanura</i>	Unpassed	EF594019	EF593977
NY03d	NY03-0199	Onondaga, NY	9/25/2003	<i>Cs. melanura</i>	Unpassed	EF594020	EF593978
NY04a	NY04-0347	Onondaga, NY	7/20/2004	<i>Cs. melanura</i>	Unpassed	EF594021	EF593979
NY04b	NY04-0390	Onondaga, NY	7/29/2004	<i>Cs. melanura</i>	Unpassed	EF594022	EF593980
NY04c	NY04-0391	Onondaga, NY	8/3/2004	<i>Cs. melanura</i>	Unpassed	EF594023	EF593981
NY04d	NY04-0384	Onondaga, NY	8/5/2004	<i>Cs. melanura</i>	Unpassed	EF594024	EF593982
NY04e	NY04-0340	Onondaga, NY	8/26/2004	<i>Cs. melanura</i>	Unpassed	EF594025	EF593983
NY04f	NY04-0401	Onondaga, NY	9/1/2004	<i>Cs. melanura</i>	Unpassed	EF594026	EF593984
NY04g	NY04-2073	Sullivan, NY	9/7/2004	<i>Haliaeetus leucocephalus</i>	Unpassed	EF594027	EF593985
NY04h	NY04-0344	Onondaga, NY	9/9/2004	<i>Cs. melanura</i>	Unpassed	EF594028	EF593986
NY04i	NY04-2727	Onondaga, NY	9/14/2004	<i>C. brachyrhynchos</i>	Unpassed	EF594029	EF593987
NY04j	NY04-0029	Ulster, NY	9/16/2004	Horse	Unpassed	EF594030	EF593988
NY04k	NY04-0037	Chemung, NY	10/10/2004	Horse	Unpassed	EF594031	EF593989
NY05a	NY05-0404	Onondaga, NY	7/14/2005	<i>Cs. Melanura</i>	v1	EF594032	EF593990
NY05b	NY05-0411	Onondaga, NY	7/26/2005	<i>Coquillettia pertubans</i>	Unpassed	EF594033	EF593991
NY05c	NY05-0443	Onondaga, NY	7/26/2005	<i>Cs. Melanura</i>	Unpassed	EF594034	EF593992
NY05d	NY05-0410	Madison, NY	8/24/2005	<i>Cs. Melanura</i>	v1	EF594035	EF593993
NY05e	NY05-2433	Onondaga, NY	9/28/2005	<i>Cyanocitta cristata</i>	Unpassed	EF594036	EF593994
NY05f	NY05-0050	Oswego, NY	10/5/2005	Horse	Unpassed	EF594037	EF593995
Ten Broeck†	EEE-Ten	Virginia	1933	Horse	Unknown	EF594038	EF593996

*Strain designations in bold were sequenced in this study. GenBank sequences from Brault et al. (1999) and Platteborze et al. (2005) (1,2).

†Passage history obtained from Brault et al. (1999) and Platteborze et al. (2005) (1,2); v, Vero cell culture; sm, suckling mouse; gp, guinea pig; ch, chicken embryo; p, unknown passages; C6/36, C636 clone of *Aedes albopictus* cell culture; bhk, baby hamster kidney cell culture; de, duck embryo cell culture; rd, human embryonal rhabdomyosarcoma cell culture; m, mosquito.

‡Information uncertain for Ten Broeck strain sequenced in this study, thought to be derived from passaged VA33 strain.

References

1. Brault AC, Powers AM, Chavez CLV, Lopez RN, Cachón MF, Gutierrez LFL, et al. Genetic and antigenic diversity among eastern equine encephalitis viruses from North, Central, and South America. *Am J Trop Med Hyg.* 1999;61:579–86. [Medline](#)
2. Platteborze PL, Kondig JP, Schoepp RJ, Wasieloski LP Jr. Comparative sequence analysis of the eastern equine encephalitis virus pathogenic strains FL91-4679 and GA97 to other North American strains. *DNA Seq.* 2005;16:308–20. [Medline](#)